

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
11 January 2001 (11.01.2001)

PCT

(10) International Publication Number
WO 01/02578 A1

(51) International Patent Classification⁷: **C12N 15/31**,
C07K 14/195, A61K 35/74, 38/16

[AU/AU]; 360 Pasir Panjang #03-11, Goldcoast Condominium, Singapore 118699 (SG).

(21) International Application Number: PCT/SG99/00128

(74) Agent: ELLA CHEONG & G. MIRANDAH; P.O. Box 0931, Raffles City, Singapore 911732 (SG).

(22) International Filing Date:

17 November 1999 (17.11.1999)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

9903146-0

2 July 1999 (02.07.1999)

SG

(71) Applicant (for all designated States except US): **INSTITUTE OF MOLECULAR AGROBIOLOGY OF 1 RESEARCH LINK** [SG/SG]; The National University of Singapore, Singapore 117604 (SG).

(81) Designated States (*national*): AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LIAN-HUI, Zhang** [AU/AU]; 360 Pasir Panjang #03-11, Goldcoast Condominium, Singapore 118699 (SG). **YIHU, Dong** [NZ/NZ]; 12 Westcoast Crescent #06-062B, Westcove Condominium, Singapore 128042 (SG). **JINLING, Xu**

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GLOBAL REGULATORS OF BACTERIAL PATHOGENIC GENES; BACTERIAL AUTOINDUCER INACTIVATION PROTEIN, AS TARGETS FOR ENGINEERING DISEASE RESISTANCE

(57) Abstract: Disclosed are an isolated nucleic acid molecule encoding a bacterial autoinducer inactivation protein, the encoded protein, and anti-bacterial uses therefor.



WO 01/02578 A1

TITLE OF THE INVENTION

**GLOBAL REGULATORS OF BACTERIAL PATHOGENIC GENES; BACTERIAL AUTOINDUCER
INACTIVATION PROTEIN, AS TARGETS FOR ENGINEERING DISEASE RESISTANCE**

CROSS-REFERENCES TO RELATED APPLICATIONS

Not applicable.

5 STATEMENT REGARDING FEDERALLY SPONSORED
RESEARCH OR DEVELOPMENT

Not applicable.

BACKGROUND OF THE INVENTION

1. Field of the Invention

10 The present invention relates to global regulators
of bacterial pathogenic genes, and their use to confer
disease resistance.

2. Description of the Related Art

15 A bibliography follows at the end of the Detailed
Description of the Invention. The listed references
are all incorporated herein by reference.

20 Cell-to-cell communication via small signal
molecules is not only of vital importance to
multi-celled living organisms such as animals and
plants, it also plays important roles in the functional
co-ordination among family members of single-celled
organisms like bacteria. Rapid progress over the last
few years has clearly established that
N-acyl-homoserine lactones, known as autoinducers

(AIs), are widely conserved signal molecules in Gram-negative bacteria. AIs were first found in marine bacteria *Vibrio* species in regulation of bioluminescence (Eberhard, et al., 1981; Cao and Meighen, 1989). In recent years, AIs have been identified in a wide range of Gram-negative bacteria. It has been found that AIs are involved in the regulation of a range of biological functions, including Ti plasmid conjugal transfer in *Agrobacterium tumefaciens* (Zhang, et al., 1993), induction of virulence genes in *Erwinia carotovora*, *Pseudomonas aeruginosa*, *Erwinia stewartii*, *Xenorhabdus nematophilus*, *Erwinia chrysanthemi*, *Pseudomonas solanaceum*, and *Xanthomonas campestris* (Jones, et al., 1993; Passador, et al., 1993; Pirhonen, et al., 1993; Pearson, et al., 1994; Beck von Bodman and Farrand, 1995; Barber, et al., 1997; Clough, et al., 1997; Costa and Loper, 1997; Dunphy, et al., 1997; Nasser, et al., 1998), regulation of antibiotics production in *Pseudomonas aureofaciens* and *Erwinia carotovora* (Pierson, et al., 1994; Costa and Loper, 1997), regulation of swarming motility in *Serratia liquifaciens* (Eberl, et al., 1996), and biofilm formation in *Pseudomonas fluorescens* and *P. aeruginosa* (Allison, et al., 1998; Davies, et al., 1998). Many more bacterial species are known to produce AIs but the biological functions related have not been established yet (Bassler, et al., 1997; Dumenyo, et al., 1998; Cha, et al., 1998).

Different bacterial species could produce different AIs. All AI derivatives share identical homoserine lactone moieties but can differ in the length and the structure of their acyl groups. The key

components in AI-mediated gene regulation systems are LuxI and LuxR type proteins. It has been established now that LuxI-type protein serves as an autoinducer synthase that utilizes acyl-ACPs and AdoMet (S-adenosylmethionine) as substrates (More, *et al.*, 1996; Schaefer, *et al.*, 1996). LuxR-type protein is proposed to be both a receptor for AIs and a AI-dependent transcriptional regulator that binds DNA immediately upstream of the *lux* promoter (Meighen, 1994; Sitnikov, *et al.*, 1995). A 20-nucleotide inverted repeat has been identified which is centered 44 nucleotides upstream of the transcription start site of the luminescence operon. This sequence called *lux* box is required for transcriptional activation by LuxR and is probably the LuxR binding site (Fuqua, *et al.*, 1994). Similar 18-bp *tra* boxes are found upstream of at least three TraR-regulated promoters, and disruption of these elements abolishes transcriptional activation by TraR (Fuqua and Winans, 1996a).

LuxR-type proteins appear to be composed of two modules (Choi and Greenberg, 1991; Hanzelka and Greenberg, 1995). Their carboxyl terminal regions contain a conserved short sequence of 19-amino acid, putative probe-type helix-turn-helix motif, predicted to be involved in binding to target promoters. A general mechanism of activation has been proposed by which the N-terminal domain of LuxR-type protein acts negatively to prevent an interaction between its C-terminal domain and the target DNA binding sites. This inhibition can be relieved by the action of an autoinducer ligand. A strong piece of evidence is that deletion of the N-terminal domain of LuxR results in constitutively active alleles of *luxR*, whereas larger

deletions that remove part of the predicted DNA binding domain abolish transcriptional activation (Choi and Greenberg, 1991). However, other members might use different mechanisms. Recent genetic studies indicate
5 that EsaR and ExpR are likely to be repressors of their target genes rather than activators. Expression of the genes that are repressed by EsaR and ExpR is increased by autoinducers (Beckvon Bodman and Farrand 1995; Throup, et al. 1995). It appears that binding of these
10 proteins to their target sites in promoter region causes repression, therefore autoinducer ligands may act to reduce binding affinity.

Evidence that the autoinducer binding site resides in the amino terminal domain of the LuxR protein has
15 been presented (Hanzelka and Greenberg, 1995). LuxR alleles that have mutated amino terminal region require higher level of this signal that does the wild type, indicating this region required for ligand interaction (Slock, et al., 1990; Shadel, et al., 1990). This
20 region (aa 79-127) and a region within the DNA-binding domain (aa 180-230) show a higher degree of conservation among LuxR and its homologs (ca 50% identity) than other parts of these polypeptides. However, the proposed protein-ligand interaction
25 between LuxR and autoinducer has not been proved yet. Analysis of merodiploid *E. coli* strains containing wild-type and mutant LuxR alleles suggested that LuxR functions as a homomultimer and that a region required for multimerization resides within amino acid residues
30 116 and 161 (Choi and Greenberg, 1992).

BRIEF SUMMARY OF THE INVENTION

In one aspect, the present invention relates to an isolated nucleic acid molecule encoding a bacterial autoinducer inactivation protein.

5 In another aspect, the present invention relates to a expression vector which comprises a nucleic acid molecule encoding a bacterial autoinducer inactivation protein, wherein the expression vector propogates in a procaryotic or eucaryotic cell.

10 In yet another aspect, the present invention relates to a cell of a procaryote or eucaryote transformed or transfected with the expression vector of the present invention.

15 In yet another aspect, the present invention relates to an isolated protein which has bacterial autoinduction inactivation activity, where the protein comprises the amino acid sequence of SEQ ID NO: 2.

20 In yet another aspect, the present invention relates to a method for increasing disease resistance in a plant or animal, which method comprises introducing into a cell of such plant or animal a nucleic acid sequence which encodes a bacterial autoinducer inactivation protein in a manner which allows said cell to express said nucleic acid sequence.

25 In yet another aspect, the present invention relates to a method of preventing or reducing bacterial damage to a plant or animal, which method comprises administering to a plant or animal in need of such prevention or reduction an effective amount of a bacterial autoinducer inactivation protein.

30 In yet another aspect, the present invention relates to a composition for reducing bacterial damage to a plant or animal, which comprises:

- a) an effective amount of a bacterial autoinducer inactivation protein; and
- b) a suitable carrier.

BRIEF DESCRIPTION OF THE DRAWINGS

5 Figure 1 shows the time course of AIs inactivation by cell extracts from *Bacillus* sp. strain 240BI. Cell extracts in 0.2 M phosphate buffer (pH 7.0) containing 100 ug total protein were added to the same buffer containing OHHL in a final concentration of 20 uM. The
10 reaction was conducted in a 1.5 ml Eppendorf centrifuge tube in a final volume of 200 microliters and incubated at 28°C. Same concentration of OHHL in the phosphate buffer was used as control. Samples were taken at
15 10-min interval till 60 min and the reaction was stopped by boiling for 3 min. The samples were centrifuged for 5 min in a bench top centrifuge at the top speed and then assayed for AIs activity as described (Zhang, 1993). Blue colony indicates the presence of AI that activates the lacZ reporter gene,
20 and white colony indicates absence of AI. Rows from left to right: 1, OHHL control without protein extract; 2 - 7, samples after 10, 20, 30, 40, 50, 60 min enzyme reaction.

 Figure 2 shows the estimation of molecular mass of
25 AIs inactivation enzyme. A 600 µl aliquot of cell extracts was added to the Centricon 30 (Amicon) and was centrifuged at a speed of 5000 x g for 30 min at 4°C. Passing fraction (550 microliters) and un-passing fraction (50 microliters) were topped up separately to
30 a final volume of 600 microliters by adding 0.2 M phosphate buffer (pH 7.0). For bioassay, different

amounts of protein samples were added to the tubes containing OHHL in a final concentration of 20 μ M. From row 1 to 6, protein samples added were 2, 4, 6, 8, 10 and 0 μ l and the final reaction volume was 20 microliters for each reaction. Plate A: Passing fraction, Plate B: un-passing fraction.

Figure 3 shows the cloning and deletion analysis of *Bacillus* SP. strain 240B1 AI inactivation region. Cosmid clone E7-R3 contains the 4.3-kb EcoRI fragment identified by restriction analysis of overlapping cosmid clones. For deletion analysis, the same fragment was cloned into cloning vector pGEM-7Zf(+) for generation of clone E7-7. The deletion subclones were produced by restriction enzyme digestion and Dnase I treatment from the clone E7-7. The location and direction of Ptac promoters in the cosmid and in the pGEM-7Zf(+) clone are indicated by arrows. AI inactivation activity of the clones is shown in the second column: +, with AI inactivation activity; -, without AI inactivation activity. Restriction enzymes: E, EcoRI; H, HindIII; Ev, EcoRV; St, StyI. The location and direction of transcription of the *aiiA* ORF is indicated by an open arrow.

Figure 4A shows the nucleotide sequence of the *aiiA* gene [SEQ ID NO:1]. The potential ribosome binding sequence and -10 promoter element are underlined and double underlined respectively. The coding portion starts at base 1. The putative factor-independent termination site is labeled by a thick underline. Figure 4B shows the predicted amino acid sequence of the *aiiA* gene product [SEQ ID NO:2].

A short peptide sequence similar to the aspartyl protease active site consensus motif is underlined.

Figure 5 shows the best match of amino acids sequence of *aiiA* gene product (AiiA) to the consensus aspartyl proteases active site motif (Asp). Symbol: X, any amino acid. A vertical line indicates perfect match.

Figure 6 shows the bioassay for Als inactivation activities in *Bacillus* sp. strain 240B1, *E. coli* clones and AIs production activity in *Erwinia carotovora* strains. Row 1, OHHL control; row 2, *Bacillus* sp. strain 240BI; row 3, *E. coli* DH5 α ; row 4, *E. coli* DH5 α (pE7-R3); row 5, *E. coli* DH5 α (pF41); row 6, *Erw. carotovora* SCG1 (pE7R3); row 7, *Erw. carotovora* SCG1 (pLAFR3); row 8, *Erw. carotovora* SCG1. In the bioassay, OHHL was added to a final concentration of 20 μ M to the samples from lines 1 to 5. No exogenous AIs were added to the samples from rows 6 to 8.

Figure 7 shows the effect of *aiiA* gene expression in *Erw. carotovora* on pathogenicity in (A), potato; (B), eggplant; (C), Chinese cabbage; (D), carrot; and (E), celery. Top: plant tissues were inoculated with *Erw. carotovora* SCG1. Bottom: plant tissues were inoculated with *Erw. carotovora* SCG1 (pE7-R3). The actively growing bacteria were centrifuged for 1 min at 3000 x g, resuspended with YEB liquid medium to OD600 = 1.3 (2×10^9 cfu/ml) which was designed as 10^0 inoculum. The 10^0 inoculum was diluted 5 and 10 times respectively to prepare $10^{-1/2}$ and 10^{-1} dilutions. The Plant tissues were inoculated by adding a 4- μ l volume of bacteria

inoculum to the freshly cut surface or a wounding site punched by a pipette tip. The inoculum concentration from the left to the right plate: 10^0 ; $10^{-1/2}$; and 10^{-1} . The inoculated plant tissues were placed in plastic
5 plates and incubated at 28°C. The photograph was taken 48 h after inoculation.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is based on the discovery that the SEQ ID NO:2 protein has the effect of reducing
10 or eliminating the activity of bacterial autoinducers (AIs). Consequently, the protein, and any nucleic acid that encodes the protein, may be used in a variety of situations where it is desired to reduce or eliminate the effect of such bacteria.

15 In one preferred aspect, the present invention provides a nucleic acid molecule which is selected from the group consisting of:

- a) a nucleic acid having the sequence of SEQ ID NO:1;
- 20 b) a nucleic acid encoding the amino acid sequence of SEQ ID NO:2; and
- c) a nucleic acid that hybridizes to a) or b) above, wherein a positive hybridization signal is observed after washing with 1 X SSC and 0.1% SDS at
25 55°C for one hour. The nucleic acid optionally further comprises a signal peptide coding region of any sequence.

The nucleic acid sequence may be used to confer bacterial resistance in plants or animals. A nucleic
30 acid that encodes a bacterial autoinducer inactivation protein can be introduced into a cell such that the

inactivation protein is expressed by the plant or animal.

The nucleic acid sequence may be used to confer resistance to diseases where the expression of pathogenic genes are regulated by autoinducers, such as the diseases caused by *Pseudomonas aeruginosa*, *Erwinia stewartii*, *Xenorhabdus nematophilus*, *Erwinia chrysanthemi*, *Pseudomonas solanaceum*, and *Xanthomonas campestris* (Passador, et al., 1993; Pirhonen, et al., 1993; Pearson, et al., 1994; Beck von Bodman and Farrand, 1995; Barber, et al., 1997; Clough, et al., 1997; Costa and Loper, 1997; Dunphy, et al., 1997; Nasser, et al., 1998). Preferably, in the agricultural setting, the sequence may be used to confer soft rot disease resistance in susceptible plants, such as potato, eggplant, Chinese cabbage, carrot and celery.

The sequence may be introduced into plant or animal cells by well-known methods. Methods for the transformation or transfection of eukaryotic cells with exogenous nucleic acid sequences include transfection, projectile bombardment, electroporation or infection by *Agrobacterium tumefaciens*. These methods are likewise familiar to the person skilled in the area of molecular biology and biotechnology and need not be explained here in detail. As pathogenic bacteria cells are confined to the intercellular area of plant tissues, it is desirable to target the AiiA protein into the intercellular spaces. Such may be accomplished by fusing a secretion signal peptide to the AiiA protein (Sato, et al., 1995; Firek, et al., 1993; Conrad and Fiedler, 1998; Borisjuk, et al., 1999). Alternatively, a plant membrane attachment motif can be incorporated into the peptide sequence of AiiA for anchoring the

AiiA enzyme in the outer surface of plant cell membrane.

The present invention provides a new strategy for engineering resistance to diseases. In particular, this strategy targets N-acyl homoserine lactone autoinducers that induce expression of pathogenic genes of many bacterial pathogens at a threshold concentration. This strategy is applicable to all plant, animal or mammal diseases where the expression of pathogenic genes of the bacterial pathogens is inducible by N-acyl homoserine lactone autoinducers.

The present invention also contemplates usage of a bacterial autoinducer inactivation protein directly to treat or prevent bacterial damage. For example, the protein may be applied directly to plants in need of such treatment or prevention. In a preferred embodiment, the protein is applied in the form of a composition which comprises an effective amount of the protein and a suitable carrier. The composition may have a wide variety of forms, including solutions, powders, emulsions, dispersions, pastes, aerosols, etc.

The bacterial autoinducer inactivation protein may also be used to treat bacterial infections in animals, including humans. In that application, an effective amount of the active ingredient is administered to an animal in need of such treatment.

For therapeutic treatment, the active ingredient may be formulated into a pharmaceutical composition, which may include, in addition to an effective amount of the active ingredient, pharmaceutically acceptable carriers, diluents, buffers, preservatives, surface active agents, and the like. Compositions may also

include one or more other active ingredients if necessary or desirable.

The pharmaceutical compositions of the present invention may be administered in a number of ways as will be apparent to one of ordinary skill in the art. Administration may be done topically, orally, by inhalation, or parenterally, for example.

Topical formulations may include ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Oral formulations include powders, granules, suspensions or solution in water or non-aqueous media, capsules or tablets, for example. Thickeners, flavorings, diluents, emulsifiers, dispersing aids or binders may be used as needed.

Parenteral formulations may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives.

The dose regimen will depend on a number of factors which may readily be determined, such as severity and responsiveness of the condition to be treated.

Aspects of the invention will now be illustrated with reference to the following non-limiting examples.

EXAMPLE 1

Bacterial isolate 240B1 was isolated from soil suspension based on its function for inactivation of N- β -oxo-hexanoyl-L-homoserine lactone (OHHL) and N- β -oxooctanoyl-L-homoserine lactone (OOHL) and N- β -oxodecanoyl-L-homoserine lactone (ODHL) (Zhang, et al., 1993). Unless otherwise stated, OHHL was used for routine bioassay. *Erwinia carotovora* strain SCG1 was isolated from Chinese cabbage leaf showing soft rot

symptoms. It has been confirmed that strain SCG1 produces AIs and elicits soft rot disease in potato and Chinese cabbage. *Escherichia coli* strain DH5 α was used as a host for DNA cloning and subcloning.

5 *Agrobacterium tumefaciens* strain NT1 (*traR*; *tra::lacZ749*) was used as an indicator in bioassay for AI activity (Piper, et al., 1993). *E. coli* strain was cultured in Luria-Bertani (LB) medium at 37°C and other strains were cultured in LB (Miller, 1972) or YEB
10 medium (per liter contains: casein hydrolysate 10 g, yeast extract 5 g, NaCl 10 g, sucrose 5 g, MgSO₄·7H₂O 0.5 g, agar 15 g, pH 7.2) at 28 °C. The minimal salts medium with mannitol and (NH₄)₂SO₄ as carbon and
15 nitrogen sources was used for bioassay of OHHL (Petit and Tempe, 1978). Appropriate antibiotics were added as indicated at the following concentrations: ampicillin, 100 µg/ml; tetracycline, 20 µg/ml and kanamycin, 50 µg/ml.

Bioassay of AIs activity

20 The qualitative and quantitative bioassay methods for determination of AIs activity has been described previously (Zhang, 1993). For determination of the AIs production ability of wild-type and genetically
25 modified *Erwinia* strains, the same bioassay procedure was used except that no OHHL was added into the bacterial culture.

Cloning and sequencing of the AiiA gene

Genomic DNA from 240B1 was digested partially with *EcoRI*. DNA fragments were ligated to the
30 dephosphorylated *EcoRI* site of cosmid vector pLAFR3 (Staskawicz, et al., 1987). Ligated DNA was packaged

with Gigapack III XL Packaging Extract (Stratagene) and transfected into *E. coli* DH5 α . Cosmid clones with OHHL inactivation activity were identified by using the bioassay method described above. Subcloning into sequencing vector pGEM-7Zf(+) was carried out by routine techniques (Sambrook, et al., 1989). Deletion analysis was carried out by using DnaseI method as described by Lin, et al. (1985). The sequencing was performed on both strands using the ABI PRISM™ dRhodamine Terminator Cycle Sequencing Ready Reaction Kit (PE Applied Biosystems). Nucleic acid sequence data and deduced amino acid sequences were analyzed with a DNASTAR™ sequence analysis software package (DNASTAR Inc.) and database searches were performed using the BLASTA search algorithm (Altschul, et al., 1990).

Genetic modification of Erwinia strain SCG1

The E7-R3 plasmid, carrying the *aiiA* gene in the cosmid vector pLAFR3, was transferred into *Erwinia* strain SCG1 by triparental mating with the helper strain RK2013 (Ditta, et al., 1980). Transconjugants were selected on the plates containing minimal medium with tetracycline and confirmed by PCR with primers specific to the *aiiA* gene.

Virulence tests

The virulence of wild-type *Erw. carotovora* strain SCG 1 and the *aiiA* gene transformant SCG1(E7-R3) was evaluated by inoculation. Four μ l of early stationary phase bacterial suspension (containing $\sim 2 \times 10^9$ cell/ml) or diluted bacteria was added to the cut surfaces or wounding sites of plant tissues. The inoculated plant

tissues were incubated in a Petri dish at 28°C overnight. The severity of soft rot was examined 48 hours after incubation.

Results

5 *Screening of bacteria that inactivate AIs*

Bacterial isolates from plant and soil samples were screened for enzymatic inactivation of AIs. A bacterial isolate 240B1, which showed a strong ability to eliminate AIs activity, was selected for further
10 study. The total protein extracts from isolate 240B1 eliminated AIs activity completely during one-hour incubation (Fig. 1), and the capacity of the protein extract to inactivate AIs was abolished by treatment with proteinase K for 1 hour or boiling for 5 min.
15 These observations indicate enzymatic inactivation of AIs by bacterial isolate 240B1. The isolate was taxonomically characterized as *Bacillus* sp., because of the following characteristics: Gram-positive, rod-shaped, catalase positive, facultatively anaerobic,
20 and 16 rRNA sequence homology to that of other *Bacillus* bacteria (data not shown).

The molecular mass of the enzyme for AIs inactivation appears to be larger than 30 kDa. Its activity was lost after passing the protein extract
25 through Centricon 30 (Amicon) but the activity was recovered in the re-suspended fraction that failed to pass the Centricon 30 (Fig. 2).

Cloning and localization of AIs inactivation region

To identify the gene encoding AIs inactivation, a
30 cosmid library was constructed with the genomic DNA of *Listera* sp. strain 240B1. Twelve hundred clones were

screened for AIs inactivation activity. Three clones showing AIs inactivating function were identified. Restriction analysis showed that the 3 clones shared one common band of 4.3-kb generated by *EcoRI* digestion.

5 The bioassay with the subclone E7-7 containing this 4.3-kb *EcoRI* fragment confirmed that this fragment encodes AIs inactivation function (Fig. 3). To identify the minimum size and the location of the AIs inactivation gene (*aiiA*), a serial of deletion clones

10 was generated by deletion from both ends of this 4.3-kb fragment with the DNaseI method (Lin, et al., 1985). The results indicated that the *aiiA* gene is contained in a 1.2 Kb fragment in clone F41 (Fig. 3).

AiiA gene encodes a novel protein

15 The 1.2-kb DNA insert in clone F41 was completely sequenced from both strands. The nucleotide sequence of *aiiA* and the predicted amino acid sequence are shown in Fig. 4. The complete sequence of the DNA insert contains 1,222 base pairs and there are 4 potential

20 in-frame open reading frames (ORF) starting from nucleotide position of 1, 42, 156 and 228 respectively (Fig. 4). Deletion analysis indicated that only the longest ORF encodes AIs inactivation function, because the clone R34, in which the 48 bp promoter region and

25 nucleotides from 1 to 13 in the longest ORF were deleted, lost AI inactivation function completely, although the remaining DNA insert was placed under the control of a functional *Ptac* promoter (Fig. 3). This is confirmed by fusing the longest ORF to the

30 glutathione S-transferase gene in the same ORF and testing for AI inactivation activity of the purified fusion protein (data not shown). This ORF contains 750

bp nucleotide and encodes a protein of 250 amino acids, with a predicted molecular mass of 28,036 daltons and an isoelectric point at 4.7, because of 19 strongly basic and 39 strongly acidic amino acids residues. The putative initiation codon is preceded at a spacing of 7 bp by a potential ribosome-binding sequence (AAGGTGG) which is complementary to the 3' end of the *E. coli* 16S rRNA. The best sequence match (TATTGT) to the consensus -10 promoter element (TATAAT) occurs 35 bp upstream of the initiation codon. A TCTT box following a T-rich region resembling the potential factor-independent termination site is found downstream of the termination codon (Brendel, 1986). The total GC content of the *aiiA* gene is 37% and GC content in the third position of the codon is 27.2%.

Database searches showed that the *aiiA* gene has no significant similarity to known sequences in the major databases (GenBank, European Molecular Biology Laboratory, Protein Information Resource, and Swiss-Prot) by FASTA and BLAST analysis at either nucleotide or peptide sequence level, suggesting that *AiiA* is a novel protein. Consensus protein motif search using the Genetics Computer Group (Madison, WI) MOTIF program showed that a short peptide sequence, "ILVDTGMPESAV" from position 47 to 58 in *AiiA*, is similar but not identical to the aspartyl protease active site signature pattern (Rawlings and Barrett, 1995) (Fig. 5).

Expression of aiiA gene in Erwinia carotovora decreases AIs releasing and attenuates virulence

The cosmid clone E7-R3 was transferred into *Erwinia carotovora* strain SCG1 by triparental mating. The pLAFR3 vector has been safely maintained in *Erwinia*

carotovora without selection pressure. The bioassay showed that the AIs released by *Erwinia carotovora* (E7-R3) was significantly reduced (Fig. 6, lane 6), while the presence of the cosmid vector pLAFR3 alone in *Erwinia carotovora* did not affect AIs production (Fig. 6, lanes 7). Data suggest that the most of AIs produced by *Erwinia carotovora* strain SCG1 was inactivated by *aiiA* gene product.

The *Erwinia carotovora* SCG1(E7-R3) that expresses *AiiA* protein failed to or caused only minor soft rot disease symptom in potato, eggplant, Chinese cabbage, carrot and celery, while its parental strain caused severe symptoms (Fig. 7A, B, C, D, E). To prevent experimental errors due to genetic variations, four colonies from *Erwinia carotovora* strain SCG1 and its *aiiA* gene transformants respectively, were randomly selected for testing AIs production and virulence on potato. Similar results were obtained in both experiments. The *Erwinia carotovora* strain SCG1 (pLAFR3) that contains the cosmid vector only caused the same level of disease severity as its parental strain *Erwinia carotovora* strain SCG1 (Fig. 7F).

Discussion

Bacterial isolate 240B I, which was identified as *Bacillus* sp., produces an enzyme that can effectively inactivate the three AIs tested, i.e., N- β -oxo-hexanoyl-L-homoserine lactone, N- β -oxo-octanoyl-L-homoserine lactone and N- β -oxo-decanoyl-L-homoserine lactone. The gene (*aiiA*) encoding the AI inactivation enzyme has been cloned and fully sequenced. Expression of the *aiiA* gene in transformed *E. coli* and pathogenic bacteria *Erwinia carotovora* confers ability for AI

inactivation and significantly reduces the AIs release from *Erwinia carotovora*. To our knowledge, it is the first protein identified capable of enzymatic inactivation of N-acyl-homoserine lactones, the autoinducers for global gene regulation in a diverse of bacteria species.

The AiiA is a novel protein. There is no significant homology to known proteins in major databases. It shares similarities to the consensus pattern of the aspartyl proteases active site (Rawlings and Barret, 1995). Aspartyl proteases, also known as acid proteases, are widely distributed in vertebrates, fungi, plants, retroviruses and some plant viruses. The aspartyl proteases from most retroviruses and some plant viruses are homodimers. The molecular mass of AiiA protein is about 28 kDa but it failed to pass a molecular sieve with a cut off size of 30 kDa, indicating a possibility that AiiA protein exists as a homodimer or homomultimer under the natural conditions. However, there is also a possibility that AiiA monomer has an irregular three-dimensional structure, which hinders it passing through the molecular sieve. Aspartyl proteases are endopeptidases and hydrolyses amide linkages of proteins. Crystallographic study has shown that the enzyme of the aspartyl protease family are bilobed molecules with the active-site cleft located between the lobes, and each lobe contributing one of the pair of aspartic acid residues that is responsible for the catalytic activity (Sielecki et al., 1991).

Erwinia carotovora is a plant pathogen that produces and secretes exoenzymes that act as virulence determinants for soft rot diseases of various plants

including potato, cabbages, tomato, chili, carrot, celery, onion, and lettuce (Kotoujansky, 1987).

Mutants that were defective in the producing N-3-(oxohexanoyl)-L-homoserine lactone were also
5 defective in synthesis of the pectinase, cellulase and protease exoenzymes. These mutants failed to induce soft rot disease in potato tubers (Jones, et al., 1993). It was found that the *expI* gene, which is homologous to *luxI* gene of *Vibrio fischeri*, encodes
10 autoinducer production in *Erwinia carotovora*. The *expI* mutant was avirulent when it was inoculated to tobacco leaf but the virulence was restored by external autoinducer addition (Pirhonen, et al., 1993). Obviously, autoinducers are a potential target for
15 genetic engineering of plant soft rot disease resistance. As an interim test and a concept proving approach, the cosmid clone containing the *aiiA* gene was introduced to *Erwinia carotovora* strain SCG1. Expression of the AiiA enzyme in *Erwinia carotovora*
20 significantly reduced the release of autoinducers, and the genetically modified *Erwinia carotovora* that expressed AiiA failed to induce any or induce only minor soft rot disease symptom on all plants tested, including potato, eggplant, Chinese cabbage, carrot and
25 celery. Our results further support the important role of autoinducers in the regulation of expression of virulence genes in *Erwinia carotovora*, and the potential of the *aiiA* gene to confer resistance to soft rot disease and other diseases in which the
30 autoinducers are involved in regulation of pathogenic gene expression.

The present invention provides a new strategy for engineering resistance to diseases. In particular,

this strategy targets N-acyl homoserine lactone autoinducers that induce expression of pathogenic genes of many bacterial pathogens at a threshold concentration. By using the above-mentioned conception-proving approach, the present invention demonstrates that reduction or elimination of autoinducers produced by pathogenic bacteria by an autoinducer inactivation enzyme significantly attenuates pathogenicity of otherwise virulent bacterial pathogen. Because the expression of pathogenic genes in pathogenic bacteria requires a threshold concentration, this AI-inactivation strategy is applicable to all plant, animal or mammal diseases where the expression of pathogenic genes of the bacterial pathogens is inducible by N-acyl homoserine lactone autoinducers.

The *aiiA* gene could also be a useful tool for investigation of the role of AIs in those bacteria where the biological functions regulated by AIs has not been established. In recent years, many more bacteria species have been shown to produce AIs (Bassler, et al., 1997; Dumenyo, et al., 1998; Cha, et al., 1998; Surette, et al., 1999). Some of them are important plant pathogens such as *Pseudomonas* and *Xanthomonas* species. The gene knock out approach based on sequence homology could be difficult. The overall levels of sequence similarity of AIs synthase and the related regulatory protein from different genera are rather low, often no higher than 28-35% identity between LuxI-type proteins and 18-25% identity for LuxR-type proteins (Fuqua et al., 1996). However, it is feasible and simple to introduce the *aiiA* gene into these

bacteria to probe the biological functions regulated by AIs.

References

- Allison, D. G. R., B. Sanjose, C. Jaspe, A. Gilbert, P. (1998). Extracellular products as mediators of the formation and detachment of *Pseudomonas fluorescens* biofilms. FEMS Microbiology Letters 167, 179- 184.
- Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990). Basic local alignment search tool. *Journal Molecular Biology* 215, 403-410.
- Barber, C. E., Tang, J. L., Feng, J. X., Pan, M. Q., Wilson, T. J., Slater, H., Dow, J. M., Williams, P., and Daniels, M. J. (1997). A novel regulatory system required for pathogenicity of *Xanthomonas campestris* is mediated by a small diffusible signal molecule. *Molecular Microbiology* 24, 555-566.
- Bassler, B. L., Greenberg, E. P. & Stevens, A. M. (1997). Cross-species induction of luminescence in the quorum-sensing bacterium *Vibrio harveyi*. *Journal of Bacteriology* 179, 4043-4045.
- Beck von Bodman, S., and Farrand, S. K. (1995). Capsular polysaccharide biosynthesis and pathogenicity in *Erwinia stewartii* require induction by an Nacylhomoserine lactone autoinducer. *Journal of Bacteriology* 177, 5000-5008.
- Borisjuk, N. V., Borisjuk, L. G., Logendra, S., Logendra, S., Petersen, F., Gleba, Y., Raskin, I. 1999. Production of recombinant proteins in plant root exudates. *Nature Biotechnology* 17, 466-469.

- Brendel, V., and Trifonov, E. N. (1984). A computer algorithm for testing potential prokaryotic terminators. *NucleicAcids Research* 12, 4411-4427.
- 5 Cao, J. G., and Meighen, E. A. (1989). Purification and structural identification of an autoinducer for the luminescence system of *Vibrio harveyi*. *Journal of Biological Chemistry* 264, 21670-21676.
- 10 Cha, C., Gao, P., Chen, Y. C., Shaw, P. D., and Farrand, S. K. (1998). Production of acyl-homoserine lactone quorum-sensing signals by gram- negative plantassociated bacteria. *Molecular and PlantMicrobe Interactions* 11, 1119-1129.
- 15 Choi S. H., Greenberg E. P. (1991). The C-terminal region of the *Vibrio fischeri* LuxR protein contains an inducer-independent *lux* gene activating domain. *Proc. Natl. Acad. Sci. USA* 88, 11115-11119.
- 20 Choi S. H, Greenberg E. P. (1992). Genetic evidence for multimerization of LuxR, the transcriptional activator of *Vibrio fischeri* luminescence. *Molec. Mar. Biol. Biotech.* 1: 408-413.
- 25 Clough, S. J., Lee, K. E., Schell, M. A., and Denny, T. P. (1997). A two-component system in *Ralstonia (Pseudomonas) solanacearum* modulates production of PhcAregulated virulence factors in response to 3-hydroxypalmitic acid methyl ester. *J Bacteriol* 179, 3639-3648.

- Collmer, A. K., N. T. (1986). The role of pectic enzymes in plant pathogenesis. *Annual Review of Phytopathology* 24, 383-409.
- 5 Conrad, U., and Fiedler, U. 1998. Compartment-specific accumulation of recombinant immunoglobulins in plant cells: an essential tool for antibody production and immunomodulation of physiological functions and pathogen activity. *Plant Molecular Biology* 38, 101-109.
- 10 Costa, J. M. & Loper, J. E. (1997). EcbI and EcbR: homologs of LuxI and LuxR affecting antibiotic and exoenzyme production by *Erwinia carotovora* subsp. *betavasculorum*. *Can J Microbiol* 43, 1164-71.
- 15 Davies, D. G., Parsek, M. R., Pearson, J. P., Iglewski, B. H., Costerton, J. W., and Greenberg, E. P. 1998. The involvement of cell-to-cell signals in the development of a bacterial biofilm. *Science* 280, 295-298.
- 20 Ditta, G., StanFleld, S., Corbin, D., and Helinski, D. R. (1980). Broad host range DNA cloning system for gram-negative bacteria: construction of a gene bank of *Rhizobium meliloti*. *Proc Natl Acad Sci USA* 77, 7347-7351.
- 25 Dumenyo, C. K. M., A. Chun, W. Chatterjee, A K. (1998). Genetic and physiological evidence for the production of N-acyl homoserine lactones by *Pseudomonas syringae* pv. *syringae* and other fluorescent plant pathogenic *Pseudomonas* species. *European Journal of Plant Pathology* 104(6). 1998. 569-582. 104, 569-582.

- Dunphy, G., Miyamoto, C., and Meighen, E. (1997). A homoserine lactone autoinducer regulates virulence of an insect-pathogenic bacterium, *Xenorhabdus nematophilus* (Enterobacteriaceae). *J Bacteriol.* 179, 5288-5291.
- Eberhard, A., Burlingame, A. L., Eberhard, C., Kenyon, G. L., Nealson, K. H., and Oppenheimer, N. J. (1981). Structural identification of autoinducer of *Phosphobacterium fischeri* luciferase. *Biochemistry* 20, 2444-2449.
- Eberl, L., Winson, M. K., Sternberg, C., Stewart, G. S. A. B., Christiansen, G., Chhabra, S. R., Bycroft, B., Williams, P., Molin, S. & Givskov, M. (1996). Involvement of N-acyl-L-homoserine lactone autoinducers in controlling the multicellular behaviour of *Serratia liquefaciens*. *Molecular Microbiology* 20, 127- 136.
- Firek, S., Draper, J., Owen M. R. L., Gandeche, A., Cockburn, B., and Whitelam, G. C. 1993. Secretion of a functional single-chain Fv protein in transgenic tobacco plants and cell suspension cultures. *Plant Molecular Biology* 23, 861-870.
- Fuqua C, Winans S. C. (1996). Conserved cis-acting promoter elements are required for density-dependent transcription of *Agrobacterium tumefaciens* conjugal transfer genes. *J. Bacteriol.* 178, 435-440.
- Fuqua W. C., Winans S. C., Greenberg E. P. (1994). Quorum sensing in bacteria: the LuxR/LuxI family of

cell density-responsive transcriptional regulators. *J. Bacteriol.* 176, 269-75.

Fuqua, C., Winans, S. C., and Greenberg, E. P. (1996).
Census and consensus in bacterial ecosystems: The
5 LuxR-LuxI family of quorum-sensing transcriptional
regulators. *Annu. Rev. Microbiol.* 50, 727-751.

Hanzelka, B. L., Greenberg, E. P. (1995). Evidence that
the N-terminal region of the *Vibrio fischeri* LuxR
protein constitutes an autoinducer-binding domain. *J.*
10 *Bacteriol.* 177, 815-817.

Jones, S. M., Yu, B., Bainton, N. J., Birdsall, M.,
Bycroft, B. W., Chhabra, S. R., Cox, A. J. R., Golby,
P., Reeves, P. J., Stephens, S., Winson, M. K.,
Salmond, G. P. C., Stewart, G. S. A. B., and Williams,
15 P. (1993). The *Lux* autoinducer regulates the production
of exoenzyme virulence determination in *Erwinia*
carotovora and *Pseudomonas aeruginosa*. *EMBO J* 12,
2477-2482.

Kotoujansky, A. (1987). Molecular genetics of
20 pathogenesis by soft-rot *Erwinias*. *Annual Review of*
Phytopathology 25, 405-430.

Lin, H. C., Lei, S. P., and Wilcox, G. (1985). An
improved DNA sequencing strategy. *Anal Biochem* 147,
114-119.

25 Meighen, E. A. (1994). Genetics of bacterial
luminescence. *Annu. Rev. Genet.* 28, 117-139.

- More, M. I., Finger, L. D., Stryker, J. L., Fuqua, C., Eberhard, A. & Winans, S. C. (1996). Enzymatic synthesis of a quorum-sensing autoinducer through use of defined substrates. *Science* 272, 1655-1658.
- 5 Nasser, W., Bouillant, M. L., Salmond, G. & Reverchon, S. (1998). Characterization of the *Erwinia chrysanthemi* *expl-expR* locus directing the synthesis of two N-acyl-homoserine lactone signal molecules. *Molecular Microbiology* 29, 1391-1405.
- 10 Passador, L., Cook, J. M., Gambello, M. J., Rust, L., Iglewski, B. H. (1993). Expression of *Pseudomonas aeruginosa* virulence genes requires cell-to-cell communication. *Science* 260, 1127- 1130.
- 15 Pearson, J. P., Gray, K. M., Passador, L., Tucker, K. D., Eberhard, A., Iglewski, B. H., and Greenberg, E. P. (1994). Structure of the autoinducer required for expression of *Pseudomonas aeruginosa* virulence genes. *Proc Natl Acad Sci USA* 91, 197-201.
- 20 Petit, A., and Tempe, J. (1978). Isolation of *Agrobacterium* Ti plasmid regulatory mutants. *Mol. Gen. Genet.* 167, 147-155.
- Pierson, L. S., 3rd, Keppenne, V. D., and Wood, D. W. (1994). Phenazine antibiotic biosynthesis in *Pseudomonas aureofaciens* 30-84 is regulated by PhzR in response to cell density. *J. Bacteriol.* 176, 3966-74.
- 25 Piper, K. R., Beck von Bodman, S., and Farrand, S. K. (1993). Conjugation factor of *Agrobacterium tumefaciens*

regulates Ti plasmid transfer by autoinduction. *Nature* 362, 448-450.

- 5 Pirhonen, M., Flego, D., Heikinheimo, R., and Palva, E. (1993). A small diffusible signal molecule is responsible for the global control of virulence and exoenzyme production in the plant pathogen *Erwinia carotovora*. *EMBO J* 12, 2467-2476.

- 10 Rawlings, N. D. B., A. J. (1995). Families of aspartic peptidases, and those of unknown catalytic mechanism. In *Methods in Enzymology*, pp. 105-180. Edited by A. J. Barrett. New York: Academic Press.

Sambrook, J. F., E. F. Maniatis, T. (1989). *Molecular Cloning*, Second edn. New York: Cold Spring Harbor Laboratory Press.

- 15 Sato, F., Koiwa, H., Sakai, Y., Yamada, Y. (1995). Synthesis and secretion of tobacco neutral PR-5 protein by transgenic tobacco and yeast. *Biochemical & Biophysical Research Communications*. 211, 909-913.

- 20 Schaefer, A. L. V., D L. Hanzelka, B L. Cronan, J E Jr. Greenberg, E P. (1996). Generation of cell-to-cell signals in quorum sensing: Acyl homoserine lactone synthase activity of a purified *Vibrio fischeri* LuxI protein. *Proceedings of the National Academy of Sciences of the United States of America* 93, 9505-9509.

- 25 Shadel, G. S., Young, R., Baldwin, T. O. (1990). Use of regulated cell lysis in a lethal genetic selection in *Escherichia coli*: identification of the

- autoinducer-binding region of the LuxR protein from *Vibrio fischeri* ATCC 7744. *J. Bacteriol.* 172, 3980-3987.
- Sielecki, A. R., Fujinaga, M., Read, R. J. & James, M. N. (1991). Refined structure of porcine pepsinogen at 1.8 Å resolution. *J Mol Biol* 219, 671-692.
- Sitnikov, D., Schineller, J. B., Baldwin, T. O. (1995). Transcriptional regulation of bioluminescence genes from *Vibrio fischeri*. *Mol. Microbiol.* 17, 801-12.
- Slock J, Kolibachuk D, Greenberg EP. (1990). Critical regions of the *Vibrio fischeri* LuxR protein defined by mutational analysis. *J. Bacteriol.* 172: 3974-3979.
- Staskawicz, B. D., D. Keen, N. T. and Napolitano, C. (1987). Molecular characterization of cloned avirulence genes from race 0 and race I of *Pseudomonas syringae* pv. *glycinea*. *Journal of Bacteriology* 169, 5789-5794.
- Surette, M. G. B., B L. (1998). Quorum sensing in *Escherichia coli* and *Salmonella typhimurium*. *Proceedings of the National Academy of Sciences of the United States of America.* 95, 7046-7050.
- Throup, J. P., Camara, M., Briggs, G. S., Winson, M. K., Chhabra, S. R., et al. (1995). Characterisation of the *yenI/yenR* locus from *Yersenia enterocolitica* mediating the synthesis of two N-acylhomoserine lactone signal molecules. *Mol. Microbiol.* 17, 345-356.
- Zhang, L.-H. (1993). Molecular biology and biochemistry of a novel conjugation factor in *Agrobacterium*.

Doctoral Dissertation, The Adelaide University,
Australia.

Zhang, L.-H., Murphy, P. J., Kerr, A., and Tate, M. E.
(1993). *Agrobacterium* conjugation and gene regulation
5 by N-acyl-L-homoserine lactones. *Nature* 362, 446-447.

CLAIMS

1. An isolated nucleic acid molecule encoding a bacterial autoinducer inactivation protein.

2. The molecule of claim 1, wherein the nucleic acid molecule is selected from the group consisting of:

a) a nucleic acid having the sequence of the coding portion of SEQ ID NO:1;

b) a nucleic acid encoding the amino acid sequence of SEQ ID NO:2; and

c) a nucleic acid that hybridizes to a) or b) above, wherein a positive hybridization signal is observed after washing with 1 X SSC and 0.1% SDS at 55°C for one hour.

3. The molecule of claim 1, which further comprises a signal peptide coding region of any sequence.

4. An expression vector which comprises the nucleic acid molecule of claim 1, wherein the expression vector propagates in a procaryotic or eucaryotic cell.

5. A cell of a procaryote or eucaryote transformed or transfected with the expression vector of claim 4.

6. An isolated protein which has bacterial autoinduction inactivation activity, where the protein comprises the amino acid sequence of SEQ ID NO: 2.

7. A method for increasing disease resistance in a plant or animal, which method comprises introducing

33

into a cell of such plant or animal a nucleic acid sequence which encodes a bacterial autoinducer inactivation protein in a manner which allows said cell to express said nucleic acid sequence.

5 8. The method of claim 7, wherein the nucleic acid sequence is selected from the group consisting of:

a) a nucleic acid having the sequence of the coding portion of SEQ ID NO:1;

10 b) a nucleic acid encoding the amino acid sequence of SEQ ID NO:2.

9. The method of claim 7 or 8, wherein the nucleic acid sequence further comprises a signal peptide coding region of any sequence.

15 10. The method of claim 7 or 8, wherein the nucleic acid sequence further comprises a membrane attachment domain-coding region of any source.

11. The method of claim 7, wherein the plant is susceptible to bacterial soft rot disease.

20 12. The method of claim 11, wherein the plant is selected from the group consisting of potato, eggplant, Chinese cabbage, carrot and celery.

25 13. The method of claim 7, wherein the plant is susceptible to a bacterial disease in which the expression of a virulence gene is regulated by an N-acyl homoserine lactone autoinducer.

14. A method of preventing or reducing bacterial damage to a plant or animal, which method comprises

administering to a plant or animal in need of such prevention or reduction an effective amount of a bacterial autoinducer inactivation protein.

15 15. The method of claim 14, wherein the protein comprises SEQ ID NO: 2.

16. A composition for preventing or reducing bacterial damage to a plant or animal, which comprises:

- 10 a) an effective amount of a bacterial autoinducer inactivation protein; and
 b) a suitable carrier.

17. The composition of claim 16, wherein the protein comprises SEQ ID NO: 2.

18. A method for screening of bacterial isolates for autoinducer inactivation activity, which comprises:

- 15 a) isolating a single colony bacterial culture from soil or plant samples;
 b) screening the culture for autoinducer inactivation activity;
 c) preparing a crude protein extract from the culture; and
20 d) confirming enzymatic inactivation of autoinducer activity by the crude protein extract.

19. A method of isolating the nucleic acid of claim 1 or claim 2, which comprises the steps of:

- 25 a) preparing a gene bank from a donor organism that contains a nucleic acid sequence coding for a protein with an autoinducer inactivation activity in a suitable host organism;
 b) screening the clones of the gene bank; and

c) isolating the clones which contain a nucleic acid coding for a protein with autoinducer inactivation activity.

20. A process as claimed in claim 19, wherein *E. coli* is used as host organism.

21. A process as claimed in claim 19, wherein the steps of preparing a gene bank, screening the clones, and isolating the clones are performed in an *E. coli* strain that does not inactivate the autoinducer.

22. A method which comprises:
a) introducing the nucleic acid sequence of claim 1 or claim 2 into a bacterial cell; and
b) screening the bacterial cell obtained from step a) for changed biological function.

23. The method of claim 22, wherein the changed biological function is a function which is lost as a result of step a).

24. The method of claim 22, wherein the changed biological function is a function which is suppressed as a result of step a).

25. The method of claim 22, wherein the changed biological function is a function which is enhanced as a result of step a).

1/8

Figure 1




Figure 2





Figure 3

Clone Activity



4 / 8

Figure 4(A)

```
..ctttactgtattgtttttattcaaaactaaatgtaaagggtggatacata   -1
ATGACAGTAAAGAAGCTTTATTTTCGTCCCAGCAGGTCGTTGTATGTTGGA   50
TCATTTCGTCTGTTAATAGTACATTAACACCAGGAGAATTATTAGACTTAC   100
CGGTTTGGTGTTATCTTTTGGGAGACTGAAGAAGGACCTATTTTAGTAGAT   150
ACAGGTATGCCAGAAAGTGCAGTTAATAATGAAGGTCTTTTAAACGGTAC   200
ATTTGTCGAAGGGCAGGTTTACCGAAAATGACTGAAGAAGATAGAATCG   250
TGAATATTTTAAAACGGGTTGGTTATGAGCCGGAAGACCTTCTTTATATT   300
ATTAGTTCTCACTTGCATTTTGATCATGCAGGAGGAAATGGCGCTTTTAT   350
AAATACACCAATCATTGTACAGCGTGCTGAATATGAGGCGGCGCAGCATA   450
GCGAAGAATATTTGAAAGAATGTATATTGCCGAATTTAAACTACAAAATC   500
ATTGAAGGTGATTATGAAGTCGTACCAGGAGTTCAATTATTGCATACACC   550
AGGCCATACTCCAGGGCATCAATCGCTATTAATTGAGACAGAAAAATCCG   600
GTCCTGTATTATTAACGATTGATGCATCGTATACGAAAGAGAATTTTGAA   650
AATGAAGTGCCATTTGCGGGATTTGATTCAGAATTAGCTTTATCTTCAAT   700
TAAACGTTTAAAAGAAGTGGTGATGAAAGAGAAGCCGATTGTTTTCTTTG   750
GACATGATATAGAGCAGGAAAGGGGATGTAAAGTGTTCCCTGAATATATA   800
```


5/8

TAGtacaaaaagtcatgagcttattcgctcatgacttttttcgtttaaag 850
Atttttttaaataagttataaaacttttttagaactatcttcatttaattg 900
Atagtacgtaagggttacatcattaggagtatcttggtgagcaatcatca 950
Cttcgttactgtgatgggtcaactacccatatgaaatattttttataagtc 1000
Ccatcctcgaaagtaatccacatatcacagtctattaaatctgaccttc 1050
Ttcatctaattgtaattttccttttttggcggtatccatactgttaatga 1100
Atgtttttaattcatctgtttttgtgagaaagatatccttttttgtttta 1150
Attgactcgacatgtatatcttttatttcttgttttcctaaaaagacagg 1200
gggctcatttgggtctcttttgagt 1222

Figure 4(B)

MTVKKLYFVPAGRCMLDHSSVNSTLTPGELLDLPVWCYLLETEEGPILVD 50
TGMPESAVNNEGLFNGTFVEGQVLPKMTEEDRIVNILKRVGYEPEDLLYI 100
ISSHLHFDHAGGNGAFINTPIIVQRAEYEAQHSSEYLKECILPNLNYKI 150
IEGDYEVVPGVQLLHTPGHTPGHQSLLIETEKSGPVLLTIDASYTEKENFE 200
NEVPFAGFDSELALSSIKRLKEVVMKEKPIVFFGHDIEQERGCKVFPEYI 250

6 / 8

Figure 5

Asp: [LIVMFGAC] - [LIVMTADN] - [LIVFSA] - D - [ST] - G - [STAV] -
 AiiA: I₄₇ L V D T G M

Asp: [STAPDENQ] - X - [LIVMFSTNC] - X - [LIVMFGTA]
 AiiA: P E S A V₅₈

Figure 6





Figure 7



INTERNATIONAL SEARCH REPORT

In International Application No

PCT/SG 99/00128

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C12N15/31 C07K14/195 A61K35/74 A61K38/16

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C12N C07K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 96 29392 A (UNISEARCH LTD ;KJELLEBERG STAFFAN (AU); STEINBERG PETER (AU); NYS) 26 September 1996 (1996-09-26) the whole document	1,3-5,7, 9-14,16, 18-22
Y	WO 98 57618 A (UNIV MONTANA RES DEV INST ;DAVIES DAVID G (US); COSTERTON JOHN WIL) 23 December 1998 (1998-12-23) the whole document	1,3-5,7, 9-14,16, 18-22
Y	US 5 591 872 A (PEARSON JAMES P ET AL) 7 January 1997 (1997-01-07) the whole document	1,3-5,7, 9-14,16, 18-22



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- "&" document member of the same patent family

Date of the actual completion of the international search

4 April 2000

Date of mailing of the international search report

12/04/2000

Name and mailing address of the ISA

European Patent Office, P.B. 6818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Hix, R

INTERNATIONAL SEARCH REPORT

In . tional Application No

PCT/SG 99/00128

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>LUO Z -Q ET AL: "THE ANTIACTIVATOR TraM INTERFERES WITH THE AUTOINDUCER-DEPENDENT BINDING OF THE QUORUM-SENSING ACTIVATOR TraR TO DNA BY INTERACTING WITH THE C-TERMINAL REGION OF THE PROTEIN" ABSTRACTS OF THE GENERAL MEETING OF THE AMERICAN SOCIETY FOR MICROBIOLOGY, 30 May 1999 (1999-05-30) - 3 June 1999 (1999-06-03), page H91 XP000884164 abstract</p>	1,3-5,7, 9-14,16, 18-22
Y	<p>SCHAEFER A L ET AL: "QUORUM SENSING IN VIBRIO FISCHERI: PROBING AUTOINDUCER-LuxR INTERACTIONS WITH AUTOINDUCER ANALOGS" JOURNAL OF BACTERIOLOGY, US, WASHINGTON, DC, vol. 178, no. 10, May 1996 (1996-05), pages 2897-2901, XP000884120 ISSN: 0021-9193 the whole document</p>	1,3-5,7, 9-14,16, 18-22
Y	<p>EBERHARD A ET AL: "ANALOGS OF THE AUTOINDUCER OF BIOLUMINESCENCE IN VIBRIO FISCHERI" ARCHIVES OF MICROBIOLOGY, vol. 146, no. 1, October 1986 (1986-10), pages 35-40, XP000884126 ISSN: 0302-8933 the whole document</p>	1,3-5,7, 9-14,16, 18-22
Y	<p>ZHU J ET AL: "ANALOGS OF THE AUTOINDUCER 3-OXOOCTANOYL-HOMOSERINE LACTONE STRONGLY INHIBIT ACTIVITY OF THE TraR PROTEIN OF AGROBACTERIUM TUMEFACIENS" JOURNAL OF BACTERIOLOGY, US, WASHINGTON, DC, vol. 180, no. 20, October 1998 (1998-10), pages 5398-5405, XP000884118 ISSN: 0021-9193 the whole document</p>	1,3-5,7, 9-14,16, 18-22
A	<p>SLOCK J ET AL: "CRITICAL REGIONS OF THE VIBRIO FISCHERI LuxR PROTEIN DEFINED BY MUTATIONAL ANALYSIS" JOURNAL OF BACTERIOLOGY, US, WASHINGTON, DC, vol. 172, no. 7, July 1990 (1990-07), pages 3974-3979, XP000884181 ISSN: 0021-9193 the whole document</p>	
	-/-	

INTERNATIONAL SEARCH REPORT

International Application No
PCT/SG 99/00128

C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>POELLINGER K A ET AL: "INTRAGENIC SUPPRESSION OF A LuxR MUTATION: CHARACTERIZATION OF AN AUTOINDUCER-INDEPENDENT LuxR" FEMS MICROBIOLOGY LETTERS, NL, AMSTERDAM, vol. 129, no. 1, 1995, pages 97-102, XP000884123 ISSN: 0378-1097 the whole document</p>	
A	<p>REVERCHON S ET AL: "INTEGRATION OF THE QUORUM-SENSING SYSTEM IN THE REGULATORY NETWORKS CONTROLLING VIRULENCE FACTOR SYNTHESIS IN ERWINIA CHRYSANTHEMI" MOLECULAR MICROBIOLOGY, GB, BLACKWELL SCIENTIFIC, OXFORD, vol. 29, no. 6, September 1998 (1998-09), pages 1407-1418, XP000884219 ISSN: 0950-382X the whole document</p>	
A	<p>CHOI S H ET AL: "THE C-TERMINAL REGION OF THE VIBRIO FISCHERI LuxR PROTEIN CONTAINS AN INDUCER-INDEPENDENT lux GENE ACTIVATING DOMAIN" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, US, NATIONAL ACADEMY OF SCIENCE. WASHINGTON, vol. 88, December 1991 (1991-12), pages 11115-11119, XP000876990 ISSN: 0027-8424 cited in the application the whole document</p>	
A	<p>HANZELKA B L ET AL: "EVIDENCE THAT THE N-TERMINAL REGION OF THE VIBRIO FISCHERI LuxR PROTEIN CONSTITUTES AN AUTOINDUCER-BINDING DOMAIN" JOURNAL OF BACTERIOLOGY, US, WASHINGTON, DC, vol. 177, no. 3, February 1995 (1995-02), pages 815-817, XP000884175 ISSN: 0021-9193 cited in the application the whole document</p>	

INTERNATIONAL SEARCH REPORT

International application No.

PCT/SG 99/00128

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 7 to 10, 14 and 15
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/SG 99/00128

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9629392 A	26-09-1996	AU 708962 B AU 4999696 A BR 9607661 A CA 2215797 A CN 1185173 A EP 0815201 A JP 11502108 T NZ 303630 A	19-08-1999 08-10-1996 16-06-1998 26-09-1996 17-06-1998 07-01-1998 23-02-1999 26-01-1998
WO 9857618 A	23-12-1998	AU 7977698 A AU 8258998 A WO 9858075 A	04-01-1999 04-01-1999 23-12-1998
US 5591872 A	07-01-1997	NONE	